

PAB750Bo01 Polyclonal Antibody

To Tumor Necrosis Factor Ligand Superfamily, Member 13 (TNFSF13)
Organism Species: Bos taurus; Bovine (Cattle)

Instruction manual

←20kDa

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

Western Blot

70

44

33

26 22

18

14

10

[PRODUCT INFORMATION]

Immunogen: TNFSF13, Cattle

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100μg Sample: Recombinant TNFSF13, Cattle

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant TNFSF13 (Ala105~Leu250) expressed in *E.coli*.

Accession No.: RPB750Bo01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- AVLTRK HKKKRSVLHL VPINITSKED SDVTEVMWQP ALQRGRGLEA QGYVVRVWDA GVYLLYSQVL FHDETFTMGQ MVSREGQGRQ ETLFRCIQSM PSNPDWAYNS CYSAGVFHLH QGDILSVVIP RARAKLSLSP HGTFLGLVKL



[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against TNFSF13. It has been selected for its ability to recognize TNFSF13 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant TNFSF13 (Ala105~Leu250) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.