

PAB499Hu02

Polyclonal Antibody to Tumor Necrosis Factor Receptor 1 (TNFR1)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

UOM: 100µl

Cross Reactivity: Mouse

Applications: WB,IHC

[IMMUNOGEN]

Immunogen: Recombinant TNFR1 (Cys248~Met428) expressed in E.coli

Accession No.: RPB499Hu02

[APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

Cloud-Clone Corp.

expiration date under appropriate storage condition.

[IDENTIFICATION]







Figure. Western Blot; Sample: Recombinant TNFRSF1A, Human.




DAB staining on IHC-P; Samples:
Human Brain Tissue; Primary Ab:
10µg/ml Rabbit Anti-Human
TNFRSF1A Antibody Second Ab:
2µg/mL HRP-Linked Caprine AntiRabbit IgG Polyclonal Antibody
(Catalog: SAA544Rb19)




DAB staining on IHC-P;
Sample: Porcine Small intestine Tissue;
Primary Ab: 10µg/ml Rabbit AntiHuman TNFRSF1A Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;
Sample: Porcine Small intestine Tissue
Primary Ab: 10µg/ml Rabbit AntiHuman TNFR1 Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)





Western Blot; Sample: Lane1: HL60
cell lysate; Lane2: A549 cell lysate;
Lane3: Rat Placenta lysate; Lane4:
 Mouse Placenta lysate
Primary Ab: 0.3µg/ml Rabbit AntiHuman TNFR1 Antibody
Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



Western Blot; Samples: Lane1: Mouse
Placenta lysate; Lane2: Rat Placenta
lysate; Lane3: Hela cell lysate;
Primary Ab: 0.5µg/ml Rabbit AntiHuman TNFR1 Antibody
Second Ab: 0.2µg/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

Coud-Clone Corp.





Knockout Varification:

Lane 1: Wild-type Hela cell lysate;

Lane 2: TNFRSF1A knockout Hela cell

lysate;

Predicted MW: 50,38,25kDa

Observed MW: 23kDa

Primary Ab: 3µg/ml Rabbit Anti-Human

TNFRSF1A Antibody

Second Ab: 0.2µg/mL HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb19)

AF488 staining on IF;

Sample: Hela cell

Primary Ab: 20µg/ml Rabbit Anti-

Human TNFR1 Antibody

Second Ab: 2µg/ml AF488-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb11)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.