

RPC016Mu01 50µg
Recombinant 17-Alpha-Hydroxylase (S17aH)
Organism Species: *Mus musculus* (Mouse)
Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[**PROPERTIES**]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Thr201~Thr507

Tags: N-terminal His Tag

Subcellular Location: Membrane

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: 100mMNaHCO₃, 500mMNaCl, pH8.3, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.6

Predicted Molecular Mass: 38.6kDa

Accurate Molecular Mass: 38kDa as determined by SDS-PAGE reducing conditions.

[**USAGE**]

Reconstitute in 100mM NaHCO₃, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[**STORAGE AND STABILITY**]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.


Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

TEGIVDVLGH SDLVDIFPWL KIFPNKNLEM IKEHTKIREK TLVEMFEKCK
EKFNSESLSS LTDILIQAKM NAENNNTGEG QDPSVFSDKH ILVTVGDIIFG
AGIETTSSVL NWILAFVLHN PEVKRKIQKE IDQYVGFST PSFNDRTHLL
MLEATIREVL RIRPVAPLLI PHKANIDSSI GEFAIPKDTL VIINLWALHH
DKNEWDQPDR FMPEFLDPT GSHLITPTPS YLPGAGPRS CIGEALARQE
LFIFMALLLQ RFDQVSDDK QLPCLVDPK VVFLIDPFKV KITVRQAWKD
AQVEVST

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.