

RPA014Mu02 10µg

Recombinant Bone Morphogenetic Protein 4 (BMP4)

Organism Species: *Mus musculus* (Mouse)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Ser22~Arg408

Tags: N-terminal His Tag

Subcellular Location: Extracellular matrix

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.0

Predicted Molecular Mass: 48.0kDa

Accurate Molecular Mass: 48kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

SHASLIPET GKKKVAEIQG HAGGRRSGQS
 HELLRDFEAT LLQMFGRLRR PPSKSAVIP DYMRDLYRLQ SGEEEEEEQS
 QGTGLEYPER PASRANTVRS FHHEEHLENI PGTSESSAFR FLFNLSSIFE
 NEVISSAELR LFREQVDQGP DWEQGFHRIN IYEVMPKPPAE MVPGHLITRL
 LDTRLVHHNV TRWETFVSP AVLRWTREKQ PNYGLAIEVT HLHQTRTHQG
 QHVRISRSLP QGSGDWAQLR PLLVTFGHDG RGHTLTRRRA KRSPKHPQR
 SRKKKNKNCRR HSLYVDFSDV GWNDWIVAPP GYQAFYCHGD CPFPLADHLN
 STNHAIQTL VNSVNSSIPK ACCVPTLSA ISMLYLDEYD KVVLKKNYQEM
 VVEGCGCR

[IDENTIFICATION]




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.