

RPF317Hu01 10 μ g
Recombinant Carboxypeptidase A4 (CPA4)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Lys55~Tyr421

Tags: N-terminal His-Tag

Tissue Specificity: Adrenal Gland, Kidney, Heart, Brain.

Subcellular Location: Secreted.

Purity: >98%

Traits: Freeze-dried powder

Buffer formulation: 100mM NaHCO₃, 500mM NaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.0

Predicted Molecular Mass: 44.9kDa

Accurate Molecular Mass: 45kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 100mM NaHCO₃, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
KSPSSF NRPVDVLVPS VSLQAFKSFL RSQGLEYAVT IEDLQALLDN
EDDEMQHNEG QERSSNNFNY GAYHSLEAIY HEMDNIAADF PDLARRVKIG
HSFENRPMYV LKFSTGKGVR RPAVWLNAGI HSREWISQAT AIWTARKIVS
DYQRDPAITS ILEKMDIFLL PVANPDGYVY TQTQNRLWRK TRSRNPGSSC
IGADPNRNWN ASFAGKGASD NPCSEVYHGP HANSEVEVKS VVDFIQKHGN
FKGFIDLHSY SQLLMYPYGY SVKKAPDAEE LDKVARLAAK ALASVSGTEY
QVGPTCTTVY PASGSSIDWA YDNGIKFAFT FELRDTGTYG FLLPANQIIP
TAEETWLGLK TIMEHVRDNL Y
```

[IDENTIFICATION]




Figure 1. SDS-PAGE