

RPB583Hu01 100µg Recombinant Cytohesin 2 (CYTH2) Organism Species: *Homo sapiens (Human) Instruction manual* 

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Coud-Clone Corp.

# [PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Met1~Arg390

Tags: N-terminal His Tag

Subcellular Location: Membrane

**Purity:** > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.8

Predicted Molecular Mass: 49.1kDa

Accurate Molecular Mass: 49kDa as determined by SDS-PAGE reducing conditions.

## [<u>USAGE</u>]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [SEQUENCE]



MEDGVYEPPD	LTPEERMELE	NIRRRKQELL	VEIQRLREEL	SEAMSEVEGL	EANEGSKTLQ
RNRKMAMGRK	KFNMDPKKGI	QFLVENELLQ	NTPEEIARFL	YKGEGLNKTA	IGDYLGEREE
LNLAVLHAFV	DLHEFTDLNL	VQALRQFLWS	FRLPGEAQKI	DRMMEAFAQR	YCLCNPGVFQ
STDTCYVLSF	AVIMLNTSLH	NPNVRDKPGL	ERFVAMNRGI	NEGGDLPEEL	LRNLYDSIRN
EPFKIPEDDG	NDLTHTFFNP	DREGWLLKLG	GGRVKTWKRR	WFILTDNCLY	YFEYTTDKEP
RGIIPLENLS	IREVDDPRKP	NCFELYIPNN	KGQLIKACKT	EADGRVVEGN	HMVYRISAPT
QEEKDEWIKS	IQAAVSVDPF	YEMLAARKKR			

# [IDENTIFICATION]

kDa 70	-
44	
33	-
26	-
22	-
18	-
14	-
10	-

## [<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.