

RPA818Ra01 100µg Recombinant Death receptor 5 (DR5) Organism Species: *Rattus norvegicus (Rat) Instruction manual* 

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Coud-Clone Corp.

# [PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Lys17~Ala336

Tags: N-terminal His Tag

Subcellular Location: Membrane

**Purity:** > 90%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% skl, 5%Trehalose.

Original Concentration: 50µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.5

Predicted Molecular Mass: 39.1kDa

Accurate Molecular Mass: 39kDa as determined by SDS-PAGE reducing conditions.

# [ <u>USAGE</u> ]

Reconstitute in  $ddH_2O$  to a concentration of 0.1-0.5 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### [SEQUENCE]

# Cond-Clone Corp.

KLAQ DRPADLQRLK QSPLECPAGQ YLSKEDGSCK ACIDGENYTS GPNVLPSCLS CRVCKEDKVI KSRCVKARNT ECECKPGSFE DKDSTEICQT CSNCTDGEDE VIPCTPKANR KCVPKNTQIP QHNLGLLIGL LASVISVVLF VAVIWKTKAW ESVCLFMARV YPGCEQDHEN TVGLSLLDAQ TSRKTNGSHH NTEPDRTQSS PLGRKLLVLA NGNNPADALK LIFERCSTEV PFNSWDRLMR HMGLTDNQIQ MVRAETQVPC EVLYQMLLKW LYQTGLGASI NHLLGALEAV GERCALEEIE DYAVKSGKFV YQNTTA

# [IDENTIFICATION]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

# [<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.