

RPA284Hu01 50µg Recombinant Heparin Cofactor II (HCII) Organism Species: *Homo sapiens (Human) Instruction manual*

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

Cond-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Asp276~Ser499

Tags: N-terminal His Tag

Subcellular Location: Endoplasmic reticulum lumen, Extracellular matrix

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 100µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.0

Predicted Molecular Mass: 29.4kDa

Accurate Molecular Mass: 31kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in ddH_2O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]



DALEN IDPATQMMIL NCIYFKGSWV NKFPVEMTHN HNFRLNEREV VKVSMMQTKG NFLAANDQEL DCDILQLEYV GGISMLIVVP HKMSGMKTLE AQLTPRVVER WQKSMTNRTR EVLLPKFKLE KNYNLVESLK LMGIRMLFDK NGNMAGISDQ RIAIDLFKHQ GTITVNEEGT QATTVTTVGF MPLSTQVRFT VDRPFLFLIY EHRTSCLLFM GRVANPSRS

[IDENTIFICATION]




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.