

RPB853Mu01 50µg

Recombinant Matrix Metalloproteinase 19 (MMP19)

Organism Species: *Mus musculus* (Mouse)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Tyr99~Asp512

Tags: N-terminal His Tag

Subcellular Location: Membrane, Secreted, Extracellular matrix

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 250µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.9

Predicted Molecular Mass: 50.3kDa

Accurate Molecular Mass: 50kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.


Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

YL
LLGHWKKNL TFRIFNVPST LSLPRVRAAL HQAFKYWSSV APLTFREVKA
GWADIRLSFH GRQSLYCSNT FDGPGKVLAH ADIPELGSIH FDKDELWTEG
TYQGVNLR II AAHEVGHALG LGHSRYTQAL MAPVYAGYQP FFKLHPDDVA
GIQALYGKRS PETRDEEEET EMLTVSPVTA KPGMPNPCS GEVDAMVLGP
RGKTYAFKGD YVWTVTDSGP GPLFQISALW EGLPGNLDAV VYSPRTRRTH
FFKGNKVWRY VDFKMSPGFP MKFN RVEPNL DAALYWPVNQ KVFLFKGSGY
WQWDELARTD LSRYPKPIKE LFTGVPDRPS AAMSWQDGQV YFFKGKEYWR
LNQQLRVAKG YPRNTTHWMH CGSQTPD TNS STGDVTPSTT DTVLGTTTST
MGSTLDIPSA TD

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.