

RPA937Hu03 50µg Recombinant Nephrin (NPHN) Organism Species: *Homo sapiens (Human) Instruction manual*

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

Cloud-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression Host: *E.coli* Residues: His705~Ser1055 Tags: N-terminal His Tag Subcellular Location: Membrane Purity: > 80% Traits: Freeze-dried powder Buffer formulation: 100mMNaHCO₃, 500mMNaCl, pH8.3, containing 1mM DTT, 0.01% SKL, 5% Trehalose . Original Concentration: 200µg/mL Applications: Positive Control; Immunogen; SDS-PAGE; WB. (May be suitable for use in other assays to be determined by the end user.) Predicted isoelectric point: 5.7 Predicted Molecular Mass: 41.7kDa

Accurate Molecular Mass: 42kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in ddH_2O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCE</u>]

Cond-Clone Corp.

HLWNVT RADDGLYQLH CQNSEGTAEA RLRLDVHYAP TIRALQDPTE VNVGGSVDIV CTVDANPILP GMFNWERLGE DEEDQSLDDM EKISRGPTGR LRIHHAKLAQ AGAYQCIVDN GVAPPARRLL RLVVRFAPQV EHPTPLTKVA AAGDSTSSAT LHCRARGVPN IVFTWTKNGV PLDLQDPRYT EHTYHQGGVH SSLLTIANVS AAQDYALFTC TATNALGSDQ TNIQLVSISR PDPPSGLKVV SLTPHSVGLE WKPGFDGGLP QRFCIRYEAL GTPGFHYVDV VPPQATTFTL TGLQPSTRYR VWLLASNALG DSGLADKGTQ LPITTPGLHQ PSGEPEDQLP TEPPS

[IDENTIFICATION]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

[<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.