

**RPJ832Hu01 50µg**

**Recombinant Phosphoinositide-3-Kinase Catalytic Delta Polypeptide (PIK3Cd)**

**Organism Species: *Homo sapiens (Human)***

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

12th Edition (Revised in Aug, 2016)

**[ PROPERTIES ]**

**Source:** Prokaryotic expression

**Host:** *E.coli*

**Residues:** Val774~Gln1044

**Tags:** N-terminal His Tag

**Subcellular Location:** Cytoplasm

**Purity:** > 97%

**Traits:** Freeze-dried powder

**Buffer formulation:** 100mMNaHCO<sub>3</sub>, 500mMNaCl, pH8.3, containing 0.01% SKL, 5% Trehalose.

**Original Concentration:** 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 8.7

**Predicted Molecular Mass:** 34.8kDa

**Accurate Molecular Mass:** 33kDa as determined by SDS-PAGE reducing conditions.

**[ USAGE ]**

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

**[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.


Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

**[ SEQUENCE ]**

VGIIFKN GDDLKQDMLT LQMIQLMDVL  
WKQEGDLDRM TPYGCLPTGD RTGLIEVVLR SDTIANIQLN KSNMAATAAF  
NKDALLNWLK SKNPGEALDR AIEEFTLSCA GYCVATYVLG IGDRHSDNIM  
IRESGQLFHI DFGHFLGNFK TKFGINRERV PFILTYDFVH VIQQGKTNNS  
EKFERFRGYC ERAYTILRRH GLLFLHLFAL MRAAGLPELS CSKDIQYLKD  
SLALGKTEEE ALKHFRVKFN EALRESWTK VNWLAHNVSK DNRQ

**[ IDENTIFICATION ]**



**[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.