

RPD327Hu01 100µg

Recombinant Proteasome 26S Subunit, ATPase 1 (PSMC1)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Met1~Leu440

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Cytoplasm

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.4

Predicted Molecular Mass: 52.8kDa

Accurate Molecular Mass: 53kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.


Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

MGQSQSGGHG PGGGKKDDKD KKKKYEPPVP TRVGKKKKKT KGPDAASKLP LVTPHTQCRL
KLLKLERIKD YLLMEEEFIR NQEQMKPLEE KQEEERSKVD DLRGTPMSVG TLEEIIDDNH
AIVSTSVGSE HYVSILSFVD KDLLEPGCSV LLNHKVHAVI GVLMDDDTDL VIVMKVEKAP
QETYADIGGL DNQIQEIKES VELPLTHPEY YEEMGIKPPK GVILYGPPGT GKTLLAKAVA
NQTSATFLRV VGSELIQKYL GDGPKLVREL FRVAEEHAPS IVFIDEIDAI GTKRYDSNSG
GEREIQRML ELLNQLDGF DSRGDVKVIMA TNRIETL DPA LIRPGRIDRK IEFPLPDEKT
KKRIFQIHTS RMTLADDVTL DDLIMAKDDL SGADIKAICT EAGLMALRER RMKVTNEDFK
KSKENVLYKK QEGTPEGLYL

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.