

RPG794Hu02 100µg

Recombinant Transcobalamin I (TCN1)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Glu24~Tyr433

Tags: Two N-terminal Tags, His-tag and SUMO-tag

Subcellular Location: Secreted

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 4.9

Predicted Molecular Mass: 59.3kDa

Accurate Molecular Mass: 60kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```

EICEVSE ENYIRLKPLL NTMIQSNYNR
GTSAVNVVLS LKLVGIQIQT LMQKMIQQIK YNVKSRLSDV SSGELALIIL
ALGVCRAEE NLIYDYHLID KLENKFQAEI ENMEAHNGTP LTNYQLSLD
VLALCLFNGN YSTAEVNHF TPENKNYYFG SQFSVDTGAM AVLALTCVKK
SLINGQIKAD EGSLKNISY TKSLVEKILS EKKENGLIGN TFSTGEAMQA
LFVSSDYNE NDWNCQQLN TVLTEISQGA FSNPNAAAQV LPALMGKTF
DINKDSSCVS ASGNFNISAD EPITVTPPDS QSYISVNYSV RINETYFTNV
TVLNGSVFLS VMEKAQKMND TIFGFTMEER SWGPYITCIQ GLCANNNDRT
YWELLSGGEP LSQGAGSYVV RGENLEVRW SKY

```

[IDENTIFICATION]




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.