

APA415Hu01 100µg

Active Elastin Microfibril Interface Located Protein 1 (EMILIN1)

Organism Species: *Homo sapiens* (Human)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Gln806~Ala1016

Tags: N-terminal His-tag

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Buffer Formulation: PBS, pH7.4, containing 0.01% SKL, 5%Trehalose .

Original Concentration: 200µg/mL

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.1

Predicted Molecular Mass: 28.8kDa

Accurate Molecular Mass: 31kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[**SEQUENCE**]

```
QLSLK DLTGPAGEAG PPGPPGLQGP PGPAGPPGSP GKDGQEGPIG
PPGPQGEQGV EGAPAAPVPQ VAFSAALSLP RSEPGTVPFD RVLLNDGGYY
DPETGVFTAP LAGRYLLSAV LTGHRHEKVE AVLSRSNQGQV ARVDSGGYEP
EGLNKPVAE SQPSPGTLGV FSLILPLQAG DTVCVDLVMG QLAHSEEPLT
IFSGALLYGD PELEHA
```

[**ACTIVITY**]

Elastin Microfibril Interface Located Protein 1 (EMILIN1) is a glycoprotein that plays an important role in the extracellular matrix (ECM). It is widely present in the extracellular matrix of arteries, lymphatic vessels and other tissues. EMILIN1 plays an important role in several biological processes, such as participating in the biosynthesis of elastic fibers (elastogenesis), maintaining vascular cell morphology and regulating cell signaling. Besides, Integrin Beta 1 (ITGb1) has been identified as an interactor of EMILIN1, thus a functional binding ELISA assay was conducted to detect the interaction of recombinant human EMILIN1 and recombinant human ITGb1. Briefly, EMILIN1 was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to ITGb1-coated microtiter wells and incubated for 1h at 37°C. Wells were washed with PBST and incubated for 1h with anti-EMILIN1 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody for 1h at 37°C, wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 μ L stop solution to the wells and read at 450/630 nm immediately. The binding activity of recombinant human EMILIN1 and recombinant human ITGb1 was shown in Figure 1, the EC50 for this effect is 6.64 μ g/mL.




Figure 1. The binding activity of recombinant human EMILIN1 and recombinant human ITGb1

[IDENTIFICATION]




Figure 2. SDS-PAGE

Sample: Active recombinant EMILIN1, Human

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.