

APD754Ra01 100µg

Active Glypican 5 (GPC5)

Organism Species: *Rattus norvegicus (Rat)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Val27~His241

Tags: N-terminal His-tag

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Buffer Formulation: PBS, pH7.4, containing 0.01% SKL, 5%Trehalose .

Original Concentration: 200µg/mL

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.3

Predicted Molecular Mass: 27.8kDa

Accurate Molecular Mass: 27kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[**SEQUENCE**]

```
VESC EEVRKLFQWR LGGAVKGLPE APRAGFDLQV CLSKNPTCCT RKMEEQYQIA AQQDLQQVLQ
TSSSTLKLLI SRNAAAFQET LETLIRQAEN YTSILFCNTY RNMALEAAAS TQEFFTDVGL YLFGADVNP
EFVNRFFDSL FPLVYNHLIN PGVTDSSLQY SECIRTARQD VSPFGNIPKR VMGQMGRSLL PGRTFLQALN
LGIEVINTTD H
```

[**ACTIVITY**]

Glypican 5 (GPC5) belongs to the glypican family of proteoglycans that are linked to the cell surface through a glycosyl-phosphatidylinositol anchor. GPC5 is expressed primarily in embryonic neurons and mesenchyme and it is implicated in a variety of physiological processes, ranging from cell proliferation to morphogenesis. Besides, Syndecan 1 (SDC1) has been identified as an interactor of GPC5, thus a functional binding ELISA assay was conducted to detect the interaction of recombinant rat GPC5 and recombinant human SDC1. Briefly, GPC5 was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to SDC1-coated microtiter wells and incubated for 1h at 37 °C. Wells were washed with PBST and incubated for 1h with anti-GPC5 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody for 1h at 37 °C, wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37 °C. Finally, add 50 μ l stop solution to the wells and read at 450/630 nm immediately. The binding activity of recombinant rat GPC5 and recombinant human SDC1 was shown in Figure 1, the EC50 for this effect is 0.6 ug/mL.




Figure 1. The binding activity of recombinant rat GPC5 and recombinant human SDC1

[**IDENTIFICATION**]




Figure 2. SDS-PAGE

Sample: Active recombinant GPC5, Rat

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.