

APF673Hu01 100µg
Active N-Myristoyltransferase 1 (NMT1)
Organism Species: *Homo sapiens* (Human)
Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Met1~Leu222

Tags: N-terminal His-tag

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Buffer Formulation: PBS, pH7.4, containing 0.01% SKL, 5%Trehalose .

Original Concentration: 750µg/mL

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.1

Predicted Molecular Mass: 28.8kDa

Accurate Molecular Mass: 34kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
MADESETAVK PPAPPLPQMM EGNNGNGHEHC SDCENEEDNS YNRGGLSPAN DTGAKKKKKK  
QKKKKKEKGSE TDSAQDQPVK MNSLPAERIQ EIQAIELFS VGQGPACTME EASKRSYQFW  
DTQPVPKLGE VVNTHGPEP DKDNIRQEPY TLPQGFTWDA LDLGDRGVLK ELYTLLNENY  
VEDDDNMFRRF DYSPEFLLWA LRPPGWLPQW HCGVRVSSR KL
```

[ACTIVITY]

N-Myristoyltransferase 1 (NMT1) is an indispensable eukaryotic enzyme that belongs to the NMT family. NMT1 can catalyze the transfer of myristoyl groups to the amino acid terminal residues of numerous proteins. This catalytic process is required for the growth and development of many eukaryotes and viruses. Furthermore, it is recognized that N-myristoylation can also occur posttranslationally on an internal glycine exposed by caspase cleavage during apoptosis. Thus a functional binding ELISA assay was conducted to detect the interaction of recombinant human NMT1 and recombinant human CASP3. Briefly, biotin-linked NMT1 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to CASP3-coated microtiter wells and incubated for 1h at 37°C. Wells were washed with PBST 3 times and




Figure 3. SDS-PAGE

Sample: Active recombinant NMT1, Human

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.