Coud-Clone Corp.

RPB287Hu01 100µg Recombinant Integrin Alpha 5 (ITGa5) Organism Species: Homo sapiens (Human) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Glu895~Ala1049	kDa	
Tags: Two N-terminal Tags, His-tag and GST-tag	94 66.2	=
Accession: P08648	45	
Host: E. coli	33	-
Subcellular Location: Membrane; Single-pass	26	-
type I membrane protein.	20	
Purity: >95%		
Endotoxin Level: <1.0EU per 1µg	14.4	
(determined by the LAL method).		
Formulation: Supplied as lyophilized form in 20mM Tris,		
150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,		
0.01% sarcosyl, 5% trehalose, and preservative.		
Predicted isoelectric point: 8.4		
Predicted Molecular Mass: 50.0kDa		
Applications: SDS-PAGE; WB; ELISA; IP.		
(May be suitable for use in other assays to be determined by the end user.)		

[<u>USAGE</u>]

Reconstitute in sterile ddH₂O.

Coud-Clone Corp.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCES</u>]

The sequence of the target protein is listed below.

EAPSRS SASSGPQILK CPEAECFRLR CELGPLHQQE SQSLQLHFRV WAKTFLQREH QPFSLQCEAV YKALKMPYRI LPRQLPQKER QVATAVQWTK AEGSYGVPLW IIILAILFGL LLLGLLIYIL YKLGFFKRSL PYGTAMEKAQ LKPPATSDA

[REFERENCES]

- 1. Birkenmeier T.M., et al. (1991) J. Biol. Chem. 266:20544-20549.
- 2. Fitzgerald L.A., et al. (1987) Biochemistry 26:8158-8165.
- 3. Barillari G., et al. (1999) Blood 94:663-672.
- 4. Barry S.T., et al. (2000) Biochem. Biophys. Res. Commun. 267:764-769.