

CSI089Mu01 Primary Mouse Periodontal Ligament Fibroblasts (PDLF) Organism Species: Mus musculus (Mouse) Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1st Edition (Revised in Jun, 2024)

### [ DESCRIPTION ]

Cell Type: Fibroblasts Synonyms: PDLCs、 PDLF Strain: Strain: BALB/c Mouse Age: 2-3 weeks Tissue Source: Periodontal Ligament tissue Disease: Normal Size: >5×10<sup>5</sup>cell/ vial

### [ PROPERTIES ]

Cell activity: >85% (Viability by Trypan Blue Exclusion).
Formulation: Frozen 1 mL or T25 flask.
Biosafety: Negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast and fungi.
Applications: For research use only. It is not approved for human or animal use, or for application in clinical diagnostic procedures.
Growth Properties: Adherent

### [ CONTENTS ]

Form & Buffer: Supplied as solution form in frozen stock solution, containing 90% FBS+10% DMSO.

### [USAGE]

Upon receiving the cells in a T-25 flask at room temperature, immediately transfer the cells to  $37^{\circ}$ C, 5% CO<sub>2</sub> incubator; the cells in vials, directly and immediately transfer the cells from dry ice to liquid nitrogen.

#### **Culture conditions:**

DMEM/F12+10%FBS+1%Fibroblast Growth Supplement+1%Penicillin-Streptomycin Solution

Temperature: 37°C

Condition: 95% air, 5% carbon dioxide

#### Cell recovery:

After receiving the cells, shake at 37°C in a water bath until completely dissolved, transfer to a 15 ml centrifuge tube, add 3-5 times complete culture solution, 1000 rpm for 5 min, discard the supernatant, and place in a T25 flask for culture.

## Cloud-Clone Corp.

#### Cell passage:

- 1. Cell passage when cell growth at 85-95%.
- 2. Discard the medium and wash with PBS 1-2 times.
- 3. Add 1 ml of Trypsin at 37°C, observe the cell under the microscope. If the cells are retracted and rounded, pat the culture flask to let the cells fall off. Stop digestion by adding 2 ml of complete medium containing 10% serum. Make it a single cell suspension.
- 4. Add the fresh medium to resuspend the cells. Unless otherwise stated, the recommended ratio of primary cells is 1/2.

### [Shipping]

Dry ice.

### [STORAGE]

Upon receiving, directly and immediately transfer the cells from dry ice to liquid nitrogen and keep the cells in liquid nitrogen until they are needed for experiments.

### [IMPORTANT NOTE]

The cell is for research use only, and we will not be responsible for any issue if the cell was used in clinical diagnostic or any other procedures.

### [Figure]




Figure 1

Figure 2




Figure 3

Figure 4

# Cloud-Clone Corp.

- Figure 1 Morphology of Mouse Periodontal Ligament Fibroblasts (Optical microscope, ×100)
- Figure 2 Morphology of Mouse Periodontal Ligament Fibroblasts (Optical microscope,×200)
- Figure 3 Immunofluorescence identification of Fibronectin specific antibody (×100)
- Figure 4 Immunofluorescence identification of Fibronectin specific antibody (×200)