

RPG476Ra01 100µg

Recombinant Histidine Decarboxylase (HDC)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Leu365~Gly643

Tags: N-terminal His Tag

Subcellular Location: Cytoplasm

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.0

Predicted Molecular Mass: 32.6kDa

Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]



LWFVIR	SFGVKNLQAH	VRHGTDMAKY	FESLVRSDPV	FEIPAERHLG	LVVFRLKGPN
CLTESVLKEI	AKTGQVFLIP	ATIQDKLIIR	FTVTSQFTTK	DDILRDWNLI	REAANLVLSQ
HCTSQPSPRA	KNLIPPPVTR	DSKDLTNGLS	LESVNEGGDD	PVQVRKIFRL	PGDSLETTMD
PFDDCFSEEA	SDTTKHKLSS	FLFSYLSVQN	KKKTMRSLSC	NSMPMSAQKS	PPPDASVKHG
			SVPSFPECSS	이 없는데 없는데 이 아이를 보고 있다면 하는데 하는데 되었다.	

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.